

---

## Homework Solutions 2 (Ryden Chapter 3)

**3** On a circle,  $r$  is constant, so  $dr = 0$ . Therefore  $dl = R \sin(r/R) d\theta$ , and the circumference is

$$C = R \sin\left(\frac{r}{R}\right) \int_0^{2\pi} d\theta = 2\pi R \sin\left(\frac{r}{R}\right)$$

The deviation from flatness is

$$\Delta C = 2\pi r - 2\pi R \sin\left(\frac{r}{R}\right) \approx 2\pi r - 2\pi R \left(\frac{r}{R} - \frac{1}{3!} \frac{r^3}{R^3}\right) = \frac{\pi}{3} \frac{r^3}{R^2}$$

where the approximate value is for  $r \ll R$ . To get  $\Delta C > 1$  m, we need  $r > 0.0053R = 3.4 \times 10^4$  m.

**4** With  $\kappa = +1$ , a sphere, the largest triangle is a great circle, with area half the sphere:  $2\pi R^2$ .

With  $\kappa = 0$ , a plane, there is no largest triangle.

With  $\kappa = -1$ , the longest possible length of the sides of a triangle are when the angles  $\alpha = \beta = \gamma \rightarrow 0$ , in which case  $\alpha + \beta + \gamma = \pi - A/R^2 = 0$ . Therefore, you can get a triangle with infinitely long sides, but the area is  $\pi R^2$ .

**5** Brute force:

$$dx = dr \sin \theta \cos \phi + r \cos \theta d\theta \cos \phi - r \sin \theta \sin \phi d\phi$$

$$dx^2 = dr^2 \sin^2 \theta \cos^2 \phi + 2 \text{ more squares and 3 cross terms}$$

$$dy = dr \sin \theta \sin \phi + r \cos \theta d\theta \sin \phi + r \sin \theta \cos \phi d\phi$$

$$dy^2 = dr^2 \sin^2 \theta \sin^2 \phi + 2 \text{ more squares and 3 cross terms}$$

$$dz = dr \cos \theta - r \sin \theta d\theta$$

$$dz^2 = dr^2 \cos^2 \theta + r^2 \sin^2 \theta d\theta^2 - 2r \sin \theta \cos \theta dr d\theta$$

Adding everything together, there are some cancellations, and we end up with

$$dx^2 + dy^2 + dz^2 = dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\phi^2)$$